The GLIB technique for genome-wide mapping of 5-hydroxymethylcytosine.

Journal: Nat Protoc

Publication Year: 2012

Authors: William A Pastor, Yun Huang, Hope R Henderson, Suneet Agarwal, Anjana Rao

PubMed link: 23018194

Funding Grants: Generation of regulatory T cells by reprogramming

Public Summary:
5-Hydroxymethylcytosine (5hmC) is a newly discovered DNA base present at detectable levels in most mammalian cell types and tissues. It is generated by Tet-enzyme-mediated oxidation of 5-methylcytosine (5mC). 5hmC is important both because of its potential role in regulating gene expression and because it may be an intermediate in DNA demethylation. Here we describe a technique termed GLIB (glucosylation, periodate oxidation and biotinylation), which combines several enzymatic and chemical modification steps to attach biotin to 5hmC. Biotin-containing genomic DNA fragments are then enriched using streptavidin beads, eluted and sequenced. GLIB is capable of quantitatively tagging and precipitating fragments containing a single 5hmC molecule. Sample preparation and GLIB can be conducted in 2-3 d.

Scientific Abstract:
5-Hydroxymethylcytosine (5hmC) is a newly discovered DNA base present at detectable levels in most mammalian cell types and tissues. It is generated by Tet-enzyme-mediated oxidation of 5-methylcytosine (5mC). 5hmC is important both because of its potential role in regulating gene expression and because it may be an intermediate in DNA demethylation. Here we describe a technique termed GLIB (glucosylation, periodate oxidation and biotinylation), which combines several enzymatic and chemical modification steps to attach biotin to 5hmC. Biotin-containing genomic DNA fragments are then enriched using streptavidin beads, eluted and sequenced. GLIB is capable of quantitatively tagging and precipitating fragments containing a single 5hmC molecule. Sample preparation and GLIB can be conducted in 2-3 d.